

Misko Dry Semolina Pasta

Environmental Product Declaration



The first EPD process certified in the food industries



Barilla
The Italian Food Company. Since 1877.



REGISTRATION NUMBER

S-P-00488

CPC CODE

2731 Uncooked pasta, not stuffed or otherwise prepared
PCR 2010:01 v. 4.01
20.09.2021

PUBLICATION DATE

2014/09/07

REVISION

5 of 2022/03/02

VALID UNTIL

2024/11/26

PROGRAMME

The International EPD® System
www.environdec.com

PROGRAMME OPERATOR

EPD International AB

This EPD has been developed in conformity to ISO 14025. An EPD should provide current information and may be updated if conditions change. The stated validity is, therefore, subject to the continued registration and publication at www.environdec.com.

1. Brand and product

THE BRAND MISKO



Leading brand of pasta in Greece, Misko was founded in 1927. It represents the local pasta tradition and is symbolized by Akakio, a monk who rides to the village market on his donkey to buy pasta. Misko became part of the Barilla Group in 1991.

Further information on www.misko.gr.

THE PLANT AND THE PROCESS

Misko dry semolina pasta is produced in Thiva plant in Greece with only durum wheat and water.

It is produced by extrusion or lamination and finally by a drying process. Every year about 30 000 tons of Misko pasta are produced.



THE PRODUCTS

Products included in the analysis are classic pasta formats (penne, spaghetti, fusilli, etc.). Shape is the only feature differentiating these products, since they are all produced using as only ingredients water and durum wheat semolina.

PRODUCT CONTENT AND NUTRITIONAL INFORMATION

The durum wheat semolina pasta concerned by this declaration is made only by durum semolina and water.

From a nutritional point of view, its main characteristics are:

NUTRITIONAL INFORMATION (per 100 g)		
Energy	kcal kJ	359 1 521
Fats <i>of which saturated</i>	grams	2.0 0.4
Carbohydrates <i>of which sugars</i>	grams	71.7 3.0
Fibres	grams	3.0
Proteins	grams	12
Sodium	grams	0.005


2. Barilla group

Passion for quality, continuous pursuit of excellent recipes and ability to combine tradition and innovation are the fundamental ingredients that have allowed a small shop of bread and pasta, opened in 1877 in Parma, to become an international player in the market of pasta, ready-to-eat sauces, baked goods and crispy breads.

The Group operates in over 100 countries through its brands, which have become the icon of excellence in the food sector, and with 30 production sites, which every year contribute to the production of over 2,099,000 tonnes of products.

With its brands - Barilla, Mulino Bianco, Pan di Stelle, Gran Cereale, Harrys, Pavesi, Wasa, Filiz, Yemina e Vesta, Misko, Voiello, Cucina Barilla, Catelli, Lancia, Tolerant and Pasta Evangelists – promotes a tasty, joyful and healthy diet, inspired by the Mediterranean diet and the Italian lifestyle.

Further information on www.barillagroup.com



Good for You, Good for the Planet

In order to make a concrete contribution to global challenges, over the years, Barilla has developed a thought enclosed in the Good for You, Good for the Planet Mission that guides, step by step and offers people good, safe, nutritionally balanced food, coming from responsible supply chains.

GOOD FOOD means taste, pleasure and a daily gesture of love for the people themselves.


HEALTHY FOOD means selected raw materials and balanced nutritional profiles to support healthy lifestyles.

FOOD SOURCED FROM RESPONSIBLE SUPPLY CHAINS means seeking the best ingredients to guarantee excellent quality, respectful of people, animals and the environment.

A commitment “from field to fork”, which has led to the development of initiatives in the various stages of the supply chain and for which all Barilla Group brands contribute through projects aiming to improve the nutritional profile of products, reinforce the sustainability of the production and supply chains and provide transparent communication to consumers.



3. Environmental performance calculations



The environmental performance of pasta was calculated using the **LCA (life cycle analysis) methodology**, including the entire production chain, starting from the cultivation of the raw material until the delivery of the finished product to the retailer.

The study was conducted following the specific product rules published for the **EPD System**: “GPC code 2371 – Uncooked pasta, not stuffed or otherwise prepared”.

The contribution to the environmental impacts brought by generic data is less than the 10% in all impact categories.

DECLARED UNIT

Data are referred to **1 kg** of product plus the related packaging. The packaging is referred to the 500 g format, reported to 1 kg of product.

SYSTEM BOUNDARIES


The processes constituting the analyzed system were organized in three successive phases, in compliance with the EPD system’s requirements.

GEOGRAPHICAL SCOPE

The geographical scope of this EPD corresponds to the distribution area of the product; concerning Misko Pasta, this is mainly Greece, where 86% of volumes are distributed; the remaining 14% volumes are mainly distributed in Eastern Europe countries, USA, Canada, Australia.



4. Durum wheat cultivation



DURUM WHEAT CULTIVATION

Durum wheat cultivation environmental performances were analysed considering the specific durum wheat origin: the Greek region.

Percentage is calculated as average purchased amounts for years 2018, 2019, 2020.

Country specific data were used for fertilizers amount, crop yields and water use. Secondary data (mainly from Ecoinvent database) were used for fertilizers production and diesel production and use.

For every involved country, yield is calculated as average of three years (2018, 2019, 2020).


The 19,5% of Greek wheat comes from agriculture that meets the standards defined by Barilla Sustainable Farming.



Barilla purchases only wheat that fulfills its high safety and quality standards. It may occur that the Italian production during one year it is not sufficient to fulfill the quantitative and qualitative demand from Barilla, that's why the percentage of grain purchased from Italy may decrease or increase from year to year.


THE EFFORTS FOR A RESPONSIBLE FARMING

Since 2010, a team of Barilla professionals has been carrying out a study designed to identify the main areas for growing durum wheat in Italy and the cultivation systems with lower environmental impact. The main results of the project have been the publication of the Handbook for sustainable cultivation of durum wheat and the development of Granoduro.net in collaboration with Horta srl, a spin-off of the Università Cattolica di Piacenza. Barilla's commitment to the future is to disseminate these practices to reduce the durum wheat supply chain's environmental impact.



With the project Sustainable Agriculture, Barilla is the winner of the 1st European CSR Award Scheme which is an initiative promoted by the European Commission with the aim to give visibility to the best practices of Corporate Social Responsibility in Europe. The project, in collaboration with HORTA Srl and Life Cycle Engineering, has allowed the definition of the guidelines for the production of durum wheat with agricultural practices with low environmental impact.

5. Milling



MILLING

Milling process environmental performances were calculated considering energy and water consumption of Volos property mill involved in the Misko dry semolina pasta production.


Primary data (2020 year) are used for water and energy consumption and waste production. Secondary data, mainly from Ecoinvent database, are used for water and energy supply.

Environmental performances related to durum wheat transport from field to mill were evaluated by means of specific hypothesis for every production area. Secondary data, mainly from Ecoinvent database, are used for transport means.

Durum wheat does not need any particular storage condition (such as refrigeration).



6. Packaging and auxiliary materials production



PRIMARY PACKAGING


Packaging environmental performances are calculated taking into account the 500 g spaghetti packaging, one of the top seller products.

The primary packaging consists in a multilayer plastic (PP) film.

Primary data (from packaging unit) are used for packaging amount and packaging materials production; data about packaging production process come from Barilla LCA database.



Since 2004, Barilla designs new packaging with the "LCA packaging design tool". It allows the assessment of the environmental impacts of the packaging solutions already during the design phase.



Packaging used for Misko pasta is designed for recycle.

PACKAGING FOR DISTRIBUTION

The packaging for transport consists in cardboard boxes (american box), used for the distribution of the product, and a plastic extensible film. Boxes are made mainly by recycled cardboard carton (pre and post consumer).


The data used have been collected by LCA database (mainly Ecoinvent).

AUXILIARY MATERIALS

Auxiliary materials environmental performances are evaluated by using primary data from plant, during 2020 year.

Secondary data (Ecoinvent) are used for environmental aspects associated to materials production.

7. Pasta production



GENERAL INFORMATION

The environmental performances related to the production process are evaluated considering primary data for energy and water consumption and the waste production of Thiva plant. Secondary data (mainly Ecoinvent) are used for the environmental aspects related to the production of energy and water.

WATER

The water consumption is evaluated using primary data. The overall value is attributed to the product using the mass allocation procedure. Data are referred to year 2020.

WASTES

The primary data are collected by the plant registrations. The overall value is attributed to the product using the mass allocation procedure. Data are referred to year 2020.

SEMOLA INPUT TRANSPORT


Environmental performances related to semolina transport from mill to plant were evaluated considering road transport (truck) from the national mill mix and the plant using 2020 primary data. Secondary data, mainly from Ecoinvent database, are used for transport means.

ELECTRICITY


Total plant electricity consumption has been divided using mass allocation. Data are referred to 2020. Environmental performances of electric energy production were evaluated considering the specific country mix (Greece) for year 2020.

NATURAL GAS

The natural gas consumption is evaluated using primary data. The overall value is attributed to the product using the mass allocation procedure. Data are referred to year 2020.



8. Distribution



DISTRIBUTION

Misko pasta is produced in the Greek plant of Thiva.

Distribution performance were calculated using the following hypotheses:


- The 86% of production is intended to the Greek market,
- The 14% is intended to export transports, addressed mainly to Albania, USA and Bulgaria.

Data are referred to year 2019.


Primary data were used for distances covered by truck and ship; secondary data (Ecoinvent database) were used for transport means.

The product does not need any particular storage condition (such as refrigeration) during distribution.

Impacts related to transport packaging end of life are calculated considering the average scenario for paper, paperboard and plastic within the most relevant distribution countries (reference: Eurostat 2017 and EPA report 2014 for USA).



9. Cooking




COOKING PHASE

The cooking phase is strictly correlated to consumer behaviour and the related impacts could be estimated taking into account the "cooking indications" that are usually provided by the company on packaging.

The impacts related to the cooking phase could be estimated considering the cooking of 1 kg of pasta and the hypothesis reported on the PCR:

- Boiling phase: 0.18 kWh per kg of water;
- Cooking phase: 0.05 kWh per minute of cooking.

10 litres of water
per kg of pasta



1 kg of pasta

1.8 kWh
per kg of pasta
(boiling phase)


0.05 kWh
per minute of cooking

Cooking environmental performances are evaluated considering the Greek residual electricity mix for local consumption. For export market the Bulgarian residual electricity mix is used.

Barilla suggested cooking method

The energy necessary for the cooking stage has a significant impact. By choosing a cooking method that uses less energy, it is possible to sensibly reduce the carbon footprint of this stage. Pasta cooking time can be divided in two parts: the time needed to boil water and the one necessary to cook pasta. Usually, after boiling water, pasta is cooked by keeping the heat on for the entire suggested cooking time, e.g. for 10 minutes (*active cooking*). However, pasta can be cooked in a more efficient way by keeping the heat on only for the first 2 minutes of cooking and then, for the remaining suggested time, the heat can be turned off while keeping the lid on the pot (*passive cooking*).

Passive cooking can reduce the carbon footprint, due to the savings of GHG emissions related to energy use, without affecting the product quality.
Considering the cooking process of a 10-minutes-cooking 80 gr portion of pasta, cooked with gas and electric stoves, these are the possible savings:




Barilla-suggested cooking method does not affect the organoleptic properties of the product but it requires more attention during the cooking phase: pay attention that pasta is completely submerged into water and mix it regularly during cooking.

*Cooking proportion is the following: 1l water x 100gr of pasta.

**The results are valid for gas and electric stove cooking.

10. Primary packaging end of life and domestic food losses



Environmental performances of packaging end of life, for local market, are calculated by means of distribution countries end of life scenarios.



















For the export markets environmental performances are elaborated considering the end of life scenarios of the most representative distribution countries (Albania, Bulgaria and USA). The average scenario is reported here. Source: Eurostat 2017, EPA report 2014.

DOMESTIC FOOD LOSSES

The impacts related to domestic food waste are estimated assuming that 2% of the pasta is not consumed and is disposed of as waste, sent to the following destinations: 50% disposal (25% landfill + 25% incineration without energy recovery), 25% composting, 25% anaerobic digestion, following the indications of the PCR document.



















11. Environmental results - Greece (for local consumption)



 USE OF RESOURCES data referred to 1 kg of product		UPSTREAM			CORE	DOWNSTREAM	TOTAL	USE STAGE		
		 Durum wheat cultivation	 Milling	 Packaging and auxiliary materials production	 Pasta production	 Distribution up to shelf		 Packaging end of life and domestic food losses	 Pasta cooking, if gas	 Pasta cooking, if electric
PRIMARY ENERGY RESOURCES - RENEWABLE data in MJ	Used as energy carrier	5,85E-02	6,70E-02	1,00E-01	2,90E-01	1,86E-03	5,17E-01	1,02E-04	4,34E-02	3,56E+00
	Used as raw materials*	0,00E+00	0,00E+00	3,73E-02	0,00E+00	0,00E+00	3,73E-02	0,00E+00	0,00E+00	0,00E+00
	Total	5,85E-02	6,70E-02	1,37E-01	2,90E-01	1,86E-03	5,54E-01	1,02E-04	4,34E-02	3,56E+00
PRIMARY ENERGY RESOURCES - NON RENEWABLE data in MJ	Used as energy carrier	4,67E+00	7,84E-01	8,03E-01	5,63E+00	1,23E+00	1,31E+01	4,49E-03	1,50E+01	4,16E+01
	Used as raw materials	0,00E+00	5,99E-05	2,15E-01	0,00E+00	0,00E+00	2,15E-01	0,00E+00	0,00E+00	0,00E+00
	Total	4,67E+00	7,84E-01	1,02E+00	5,63E+00	1,23E+00	1,33E+01	4,49E-03	1,50E+01	4,16E+01
Secondary Material (g)		0,00E+00	0,00E+00	1,96E+01	0,00E+00	0,00E+00	1,96E+01	0,00E+00	0,00E+00	0,00E+00
Renewable secondary fuels (MJ. net calorific power)		0,00E+00	0,00E+00	1,17E-02	0,00E+00	0,00E+00	1,17E-02	0,00E+00	0,00E+00	0,00E+00
Non-renewable secondary fuels (MJ. net calorific power)		0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00
Net use of fresh water (liters)		2,01E+00	2,15E-01	3,66E-01	1,27E+00	5,25E-02	3,91E+00	7,91E-03	1,09E+01	1,89E+01
 OUTPUT FLOWS data referred to 1 kg of product		UPSTREAM			CORE	DOWNSTREAM	TOTAL	USE STAGE		
		 Durum wheat cultivation	 Milling	 Packaging and auxiliary materials production	 Pasta production	 Distribution up to shelf		 Packaging end of life and domestic food losses	 Pasta cooking, if gas	 Pasta cooking, if electric
Waste to animal feed or similar (g)		0,00E+00	0,00E+00	0,00E+00	2,77E+01	0,00E+00	2,77E+01	0,00E+00	0,00E+00	0,00E+00
Components for reuse (g)		0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00
Materials for recycling (g)		0,00E+00	7,90E-01	3,94E+00	9,36E+00	1,76E+01	3,17E+01	7,83E+00	0,00E+00	0,00E+00
Materials for energy recovery (g)		0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00	5,00E+00	0,00E+00	0,00E+00
Exported energy. electricity (MJ)		0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00
Exported energy. thermal (MJ)		0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00

Secondary energy resources and recovered energy flows do not show relevant contributions.


*The biomasses transformed into the product are not considered.

 POTENTIAL ENVIRONMENTAL IMPACTS data referred to 1 kg of product		UPSTREAM			CORE	DOWNSTREAM	TOTAL	USE STAGE		
		 Durum wheat cultivation	 Milling	 Packaging and auxiliary materials production	 Pasta production	 Distribution up to shelf		 Packaging end of life and domestic food losses	 Pasta cooking, if gas	 Pasta cooking, if electric
GLOBAL WARMING POTENTIAL - GWP (g CO ₂ eq)	Fossil	4,83E+02	4,31E+01	3,86E+01	3,26E+02	8,81E+01	9,79E+02	3,86E+00	8,86E+02	2,29E+03
	Biogenic	8,05E-02	1,36E-01	7,83E-02	5,78E-01	3,96E-03	8,77E-01	2,89E+00	3,76E-01	5,66E-01
	Land use and land transformation	3,01E-01	2,46E-03	6,87E-01	7,44E-03	7,18E-04	9,99E-01	1,08E-04	3,92E-02	9,76E-02
	Total	4,84E+02	4,32E+01	3,94E+01	3,27E+02	8,81E+01	9,81E+02	6,75E+00	8,86E+02	2,29E+03
Acidification Potential - g SO ₂ eq		1,25E+01	1,55E-01	1,29E-01	9,06E-01	4,11E-01	1,41E+01	2,78E-03	1,27E+00	8,31E+00
Eutrophication Potential - g PO ₄ ³⁻ eq		7,38E+00	2,01E-02	3,54E-02	1,13E-01	6,02E-02	7,60E+00	3,42E-03	2,32E-01	1,18E+00
Photochemical Oxidant Formation Potential - gNMVOC eq		2,09E+00	8,42E-02	1,26E-01	6,18E-01	4,99E-01	3,42E+00	4,22E-03	1,17E+00	4,51E+00
Abiotic Depletion Potential - Elements g Sb eq		6,68E-04	3,37E-07	1,27E-05	3,04E-06	3,79E-06	6,88E-04	3,61E-08	7,10E-06	2,44E-05
Abiotic Depletion Potential - Fossil fuels - MJ. net calorific value		4,53E+00	7,44E-01	9,51E-01	5,27E+00	1,23E+00	1,27E+01	4,33E-03	1,49E+01	3,95E+01
Water scarcity potential. m ³ eq		2,46E-01	5,58E-03	1,42E-02	5,82E-02	-2,63E-04	3,23E-01	2,08E-04	2,34E-01	1,71E-01
 WASTE PRODUCTION data referred to 1 kg of product		UPSTREAM			CORE	DOWNSTREAM	TOTAL	USE STAGE		
		 Durum wheat cultivation	 Milling	 Packaging and auxiliary materials production	 Pasta production	 Distribution up to shelf		 Packaging end of life and domestic food losses	 Pasta cooking, if gas	 Pasta cooking, if electric
Hazardous waste disposed (g)*		8,73E-05	0,00E+00	6,26E-04	0,00E+00	0,00E+00	7,13E-04	0,00E+00	0,00E+00	0,00E+00
Non-Hazardous waste disposed (g)*		1,38E+00	0,00E+00	2,59E+00	0,00E+00	0,00E+00	3,97E+00	0,00E+00	0,00E+00	0,00E+00
Radioactive waste disposed (g)		1,91E+02	5,14E+01	6,38E+01	2,35E+02	4,01E+01	5,81E+02	2,73E-01	6,50E+01	2,75E+03

The biogenic contribution to Global Warming Potential refers only to biogenic methane.
The contribution given by biogenic CO₂ is equal to zero, since the absorbed amount is equal to the emitted biogenic CO₂ within the reference 100 years period.



















*Zero values indicate that – even if some waste are produced and disposed – their impact is evaluated within the system boundaries.

PRODUCT ENVIRONMENTAL PERFORMANCES





















Cooking environmental performances are referred to pasta consumption in Greece.

12. Environmental results - Greece (for export destination)

 USE OF RESOURCES data referred to 1 kg of product		UPSTREAM			CORE	DOWNSTREAM	TOTAL	USE STAGE		
		 Durum wheat cultivation	 Milling	 Packaging and auxiliary materials production	 Pasta production	 Distribution up to shelf		 Packaging end of life and domestic food losses	 Pasta cooking, if gas	 Pasta cooking, if electric
PRIMARY ENERGY RESOURCES - RENEWABLE data in MJ	Used as energy carrier	5,85E-02	6,70E-02	1,00E-01	2,90E-01	3,51E-03	5,19E-01	1,04E-04	5,23E-02	3,62E+00
	Used as raw materials*	0,00E+00	0,00E+00	3,73E-02	0,00E+00	0,00E+00	3,73E-02	0,00E+00	0,00E+00	0,00E+00
	Total	5,85E-02	6,70E-02	1,37E-01	2,90E-01	3,51E-03	5,56E-01	1,04E-04	5,23E-02	3,62E+00
PRIMARY ENERGY RESOURCES - NON RENEWABLE data in MJ	Used as energy carrier	4,67E+00	7,84E-01	8,03E-01	5,63E+00	2,32E+00	1,42E+01	4,55E-03	1,33E+01	5,96E+01
	Used as raw materials	0,00E+00	5,99E-05	2,15E-01	0,00E+00	0,00E+00	2,15E-01	0,00E+00	0,00E+00	0,00E+00
	Total	4,67E+00	7,84E-01	1,02E+00	5,63E+00	2,32E+00	1,44E+01	4,55E-03	1,33E+01	5,96E+01
Secondary Material (g)		0,00E+00	0,00E+00	1,96E+01	0,00E+00	0,00E+00	1,96E+01	0,00E+00	0,00E+00	0,00E+00
Renewable secondary fuels (MJ. net calorific power)		0,00E+00	0,00E+00	1,17E-02	0,00E+00	0,00E+00	1,17E-02	0,00E+00	0,00E+00	0,00E+00
Non-renewable secondary fuels (MJ. net calorific power)		0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00
Net use of fresh water (liters)		2,01E+00	2,15E-01	3,66E-01	1,27E+00	9,82E-02	3,96E+00	7,87E-03	1,09E+01	2,19E+01
 OUTPUT FLOWS data referred to 1 kg of product		UPSTREAM			CORE	DOWNSTREAM	TOTAL	USE STAGE		
		 Durum wheat cultivation	 Milling	 Packaging and auxiliary materials production	 Pasta production	 Distribution up to shelf		 Packaging end of life and domestic food losses	 Pasta cooking, if gas	 Pasta cooking, if electric
Waste to animal feed or similar (g)		0,00E+00	0,00E+00	0,00E+00	2,77E+01	0,00E+00	2,77E+01	0,00E+00	0,00E+00	0,00E+00
Components for reuse (g)		0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00
Materials for recycling (g)		0,00E+00	7,90E-01	3,94E+00	9,36E+00	1,21E+01	2,62E+01	6,97E+00	0,00E+00	0,00E+00
Materials for energy recovery (g)		0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00	5,00E+00	0,00E+00	0,00E+00
Exported energy. electricity (MJ)		0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00
Exported energy. thermal (MJ)		0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00

Secondary energy resources and recovered energy flows do not show relevant contributions.


*The biomasses transformed into the product are not considered.

 POTENTIAL ENVIRONMENTAL IMPACTS data referred to 1 kg of product		UPSTREAM			CORE	DOWNSTREAM	TOTAL	USE STAGE		
		 Durum wheat cultivation	 Milling	 Packaging and auxiliary materials production	 Pasta production	 Distribution up to shelf		 Packaging end of life and domestic food losses	 Pasta cooking, if gas	 Pasta cooking, if electric
GLOBAL WARMING POTENTIAL - GWP (g CO ₂ eq)	Fossil	4,83E+02	4,31E+01	3,86E+01	3,26E+02	1,67E+02	1,06E+03	3,72E+00	7,23E+02	2,39E+03
	Biogenic	8,05E-02	1,36E-01	7,83E-02	5,78E-01	8,72E+00	9,59E+00	2,90E+00	2,62E-01	5,42E-01
	Land use and land transformation	3,01E-01	2,46E-03	6,87E-01	7,44E-03	1,52E-03	1,00E+00	1,10E-04	3,55E-02	2,52E-01
	Total	4,84E+02	4,32E+01	3,94E+01	3,27E+02	1,76E+02	1,07E+03	6,62E+00	7,23E+02	2,39E+03
Acidification Potential - g SO ₂ eq		1,25E+01	1,55E-01	1,29E-01	9,06E-01	1,34E+00	1,50E+01	2,80E-03	6,11E-01	1,29E+01
Eutrophication Potential - g PO ₄ ³⁻ eq		7,38E+00	2,01E-02	3,54E-02	1,13E-01	1,49E-01	7,69E+00	3,44E-03	1,85E-01	1,38E+00
Photochemical Oxidant Formation Potential - gNMVOC eq		2,09E+00	8,42E-02	1,26E-01	6,18E-01	1,23E+00	4,15E+00	4,28E-03	6,31E-01	6,90E+00
Abiotic Depletion Potential - Elements g Sb eq		6,68E-04	3,37E-07	1,27E-05	3,04E-06	6,20E-06	6,90E-04	3,55E-08	7,70E-06	9,17E-05
Abiotic Depletion Potential - Fossil fuels - MJ. net calorific value		4,53E+00	7,44E-01	9,51E-01	5,27E+00	2,31E+00	1,38E+01	4,39E-03	1,32E+01	3,86E+01
Water scarcity potential. m ³ eq		2,46E-01	5,58E-03	1,42E-02	5,82E-02	-4,84E-04	3,23E-01	2,08E-04	1,16E-01	4,25E-01
 WASTE PRODUCTION data referred to 1 kg of product		 Durum wheat cultivation	 Milling	 Packaging and auxiliary materials production	 Pasta production	 Distribution up to shelf	TOTAL	 Packaging end of life and domestic food losses	 Pasta cooking, if gas	 Pasta cooking, if electric
Hazardous waste disposed (g)*		8,73E-05	0,00E+00	6,26E-04	0,00E+00	0,00E+00	7,13E-04	0,00E+00	0,00E+00	0,00E+00
Non-Hazardous waste disposed (g)*		1,38E+00	0,00E+00	2,59E+00	0,00E+00	0,00E+00	3,97E+00	0,00E+00	0,00E+00	0,00E+00
Radioactive waste disposed (g)		1,91E+02	5,14E+01	6,38E+01	2,35E+02	7,49E+01	6,16E+02	2,78E-01	1,68E+02	2,64E+04

The biogenic contribution to Global Warming Potential refers only to biogenic methane.
The contribution given by biogenic CO₂ is equal to zero, since the absorbed amount is equal to the emitted biogenic CO₂ within the reference 100 years period.

*Zero values indicate that – even if some waste are produced and disposed – their impact is evaluated within the system boundaries.

PRODUCT ENVIRONMENTAL PERFORMANCES



Cooking environmental performances are referred to the export country with the highest distributed volumes (Albania).

13. Difference versus previous versions of the EPD

The differences versus previous EPD versions are due mainly to: updated yields for durum wheat cultivation, new input of environmental performances of plant and mills auxiliary materials, updated emission factors

for the specific energy mixes. Finally the Product Environmental Performance section has been modified with the substitution of Virtual Water Content with Water Scarcity indicator.

14. Additional information

REFERENCES

- International EPD Consortium, General Programme Instructions (EPD), ver. 3.01 of 18/09/2019;
- WWF, Global Footprint Network, Zoological Society of London, Living Planet Report 2008, WWF (2008);
- Arjen Y. Hoekstra, Ashok K. Chapagain, Maite M. Aldaya, Mesfin M. Mekonnen; Water Footprint The Water Footprint Manual 2011, Waterfootprint Network;
- PCR 2010:01; CPC 2371 - PCR for uncooked pasta, not stuffed or otherwise prepared; ver. 4.01 of 2021-09-20;
- Eurostat database for waste management, latest version (2018)



Environmental declarations published within the same product category, though originating from different programs, may not be comparable. This declaration and further information in regards are available at www.environdec.com

As EPD owner, Barilla has the sole ownership, liability and responsibility for the EPD.

EPD PROCESS CERTIFICATION

Product category Rules (PCR) review conducted by:
Technical Committee of the International EPD® system.
Chair Filippo Sessa
Contact via info@environdec.com

Program operator:
EPD International AB
Box 210 60, SE-100 31 Stockholm, Sweden
info@environdec.com



EPD PROCESS CERTIFICATION

Independent verification of the declaration and data, according to ISO 14025:

- ☒ EPD process verification
☐ EPD verification- Third party verifier

PROCESS INTERNAL VERIFICATION

Procedure for follow-up of data during EPD validity involves third part verifier:

- ☐ Yes
☒ No

Third party verifier: **Bureau Veritas Certification Sweden AB**, Accredited by: **SWEDAC**



Process internal verifier: **Ugo Pretato**, Approved by: **The International EPD® System**

STUDIO FIESCHI
& SOCI

CONTACTS

Barilla G. e R. Fratelli- Società per Azioni, via Mantova 166, 43122, Parma, Italy. www.barillagroup.com

For additional information relative to the activities of the Barilla Group or in regards to this environmental declaration, please contact:

Laura Marchelli - laura.marchelli@barilla.com



Technical support and graphic design: **Life Cycle Engineering SpA** - Italy www.lcengineering.eu



15. Glossary

ECOLOGICAL FOOTPRINT

The ecological footprint measures the area of biologically productive land and water required to provide the resources used and absorb the carbon dioxide waste generated along the entire life cycle. It is measured in standard units called global hectares (gha).

www.globalfootprint.org

CARBON FOOTPRINT

A product carbon footprint is the total amount of greenhouse gases produced along the entire life cycle. It is expressed in equivalent mass of carbon dioxide (CO₂-eq). In agriculture a significant contribution is given by the emission of nitrous oxide (N₂O) due to the fertilizers use. It is also known as Global Warming Potential (GWP).

www.ipcc.ch

WATER SCARCITY

Water scarcity measures the available water remaining per unit of surface in a given watershed relative to the world average, after human and aquatic ecosystem demands have been met. This method builds on the assumption that the potential to deprive another user of water is directly proportional to the amount of water consumed and inversely proportional to the available water remaining per unit of surface and time in a region (watershed).

www.wulca-waterlca.org

ACIDIFICATION (AP)

It is a phenomenon for which precipitation is unusually acidic, meaning that it has substandard levels of pH. It can have harmful effects on plants, aquatic animals and infrastructure. Acid rain is caused by emissions of SO₂, NO_x and NH₃. The acidification potential is measured in mass of sulphur dioxide equivalent (SO₂-eq).

EUTROPHICATION (EP)

It is an abnormal proliferation of vegetation in the aquatic ecosystems caused by the addition of nutrients into rivers, lakes or ocean, which determines a lack of oxygen. The eutrophication potential is mainly influenced by emission into water of phosphates and nitrates. It is expressed in mass of PO₄⁻ equivalent.

PHOTOCHEMICAL OXIDANT FORMATION POTENTIAL (POFP)

Production of compounds that, under the light effect, are able to promote an oxidation reaction leading to ozone production in the troposphere. The indicator is mainly influenced by VOCs (Volatile organic compounds) is usually expressed in mass of ethylene equivalent (g NMVOC - equivalent).